

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)**ScienceDirect**

Taiwanese Journal of Obstetrics &amp; Gynecology 52 (2013) 607–610

[www.tjog-online.com](http://www.tjog-online.com)

## Research Letter

# Pfeiffer syndrome with *FGFR2* W290C mutation perinatally presenting extreme proptosis

Chih-Ping Chen<sup>a,b,c,d,e,f,g,\*</sup>, Hsu-Kuang Huang<sup>h</sup>, Yu-Peng Liu<sup>i,j</sup>, Schu-Rern Chern<sup>b</sup>,  
Jun-Wei Su<sup>a,k</sup>, Wayseen Wang<sup>b,l</sup>

<sup>a</sup> Department of Obstetrics and Gynecology, Mackay Memorial Hospital, Taipei, Taiwan<sup>b</sup> Department of Medical Research, Mackay Memorial Hospital, Taipei, Taiwan<sup>c</sup> Department of Medicine, Mackay Medical College, New Taipei City, Taiwan<sup>d</sup> Department of Biotechnology, Asia University, Taichung, Taiwan<sup>e</sup> School of Chinese Medicine, College of Chinese Medicine, China Medical University, Taichung, Taiwan<sup>f</sup> Institute of Clinical and Community Health Nursing, National Yang-Ming University, Taipei, Taiwan<sup>g</sup> Department of Obstetrics and Gynecology, School of Medicine, National Yang-Ming University, Taipei, Taiwan<sup>h</sup> Department of Obstetrics and Gynecology, Taiwan Adventist Hospital, Taipei, Taiwan<sup>i</sup> Department of Radiology, Mackay Memorial Hospital Hsinchu Branch, Hsinchu, Taiwan<sup>j</sup> Mackay Junior College of Medicine, Nursing and Management, Taipei, Taiwan<sup>k</sup> Department of Obstetrics and Gynecology, China Medical University Hospital, Taichung, Taiwan<sup>l</sup> Department of Bioengineering, Tatung University, Taipei, Taiwan

Accepted 24 January 2013

The male proband was the first child of a healthy unrelated couple. The mother was 31 years old. There was no family history of congenital malformations. The pregnancy was uncomplicated except that big eyes, especially the right, and prominent forehead were noted during prenatal ultrasound examination. He was born at 37 weeks by cesarean section due to breech presentation. Birth weight was 3500 g, length 51 cm, and head circumference 37 cm. After birth, he was found to have scaphocephaly, broad big toes and thumbs, midface hypoplasia, hypertelorism, low-set ears, and severe proptosis (Fig. 1). Radiographs showed radio-ulnar-humeral synostosis, multisynostoses of sagittal and coronal sutures, brachycephaly, and ventriculomegaly (Fig. 2). A diagnosis of type 3 Pfeiffer syndrome was made. Molecular analysis of peripheral blood revealed a heterozygous c.870G>C, TGG>TGC transversion, leading to a p.Trp290Cys (W290C) mutation in the *FGFR2* gene (Fig. 3).

Pfeiffer syndrome (OMIM 101600) is an autosomal dominant disorder characterized by craniosynostosis, broad and deviated thumbs, big toes, and partial syndactyly on hands and feet, and affects about one in 100,000 individuals [1]. Three subtypes of Pfeiffer syndrome have been identified [2]. Types 2 and 3 are more common and severe than type 1. Type 1 is

classic Pfeiffer syndrome with mild manifestations of brachycephaly, midface hypoplasia, and abnormalities of the digits, and has normal intelligence, a good outcome, and a familial history of inheritance. Types 2 and 3 are associated with severe neurological compromise, poor prognosis, early death, and sporadic occurrence. Type 2 consists of cloverleaf skull, extreme proptosis, digital abnormalities, ankylosis of elbows, and developmental delay. Type 3 is similar to type 2 but without cloverleaf skull. Type 1 Pfeiffer syndrome is caused by mutations in *FGFR1* (5%) or *FGFR2* (95%), whereas types 2 and 3 Pfeiffer syndrome are caused by mutations in *FGFR2* (100%) only [3]. Hearing loss and hydrocephalus can be seen in type 1, and choanal stenosis/atresia,



Fig. 1. Craniofacial appearance and extreme proptosis at birth.

\* Corresponding author. Department of Obstetrics and Gynecology, Mackay Memorial Hospital, 92, Section 2, Chung-Shan North Road, Taipei, Taiwan.

E-mail address: [cpc\\_mmh@yahoo.com](mailto:cpc_mmh@yahoo.com) (C.-P. Chen).

laryngotracheal abnormalities, cleft palate, hydrocephalus, seizures, sacrococcygeal eversion/appendage, and early death can be seen in types 2 and 3 [3–7].

Prenatal diagnosis of craniosynostosis associated with digital abnormalities should include a differential diagnosis of Apert syndrome and Pfeiffer syndrome, both of which can be caused by *FGFR2* mutations. Apert syndrome is usually associated with *FGFR2* S252W and P253R mutations, and severe syndactyly of hands and feet, but no cloverleaf skull or

proptosis. Pfeiffer syndrome is usually associated with *FGFR2* W290C, Y340C, C342R, and S351C mutations, and can be associated with cloverleaf skull, proptosis, and broad great toes and thumbs [5,8].

The present case was associated with the *FGFR2* W290C mutation and type 3 Pfeiffer syndrome. To date, at least 16 cases of Pfeiffer syndrome with the *FGFR2* W290C mutation have been reported [5,9–17]. All were associated with type 2 or 3 Pfeiffer syndrome and had severe phenotypic features. The




Fig. 2. Radiographs show (A) radio–ulnar–humeral synostosis (white arrows); (B) multisynostoses of sagittal and coronal sutures (black arrows), shallow orbits, and brachycephaly; and (C) ventriculomegaly.

*FGFR2* genotype: c.870G→C/WT  
Proband




Fig. 3. Molecular analysis of the *FGFR2* gene shows a heterozygous c.870G>C, TGG>TGC transversion, leading to a p.Trp290Cys (W290C) mutation in the proband. WT = wild type.

*FGFR2* mutations of W290C (p.Trp290Cys), Y340C (p.Tyr340Cys), C342R (p.Cys342Arg), and S351C (p.Ser351-Cys) have been associated with severe phenotypic features of Pfeiffer syndrome such as cloverleaf skull, extreme proptosis, midface hypoplasia, hydrocephalus, tracheal sleeve, Chiari malformation, radio-ulnar-humeral synostosis, and early infant death [3–5,13,14,16,18–21]. W290C (p.Trp290Cys) creating cysteine residue has been noted to cause severe Pfeiffer syndrome, whereas W290G (p.Trp290Gly) and W290R (p.Trp290Arg) cause mild Crouzon syndrome; the same is true for Y340C (p.Tyr340Cys) versus Y340H (p.Tyr340His) [16]. Y340C causes severe Pfeiffer syndrome, whereas Y340H causes mild Crouzon syndrome.

Ocular manifestations play an essential role in the diagnosis and management of Pfeiffer syndrome. The present case presented ocular manifestations of shallow orbits, proptosis, and hypertelorism. Types 2 and 3 Pfeiffer syndrome tend to have more severe ocular proptosis secondary to a shallow orbit, an increased incidence of spontaneous globe subluxation, and a less favorable prognosis. Okajima et al [22] demonstrated that *FGFR2* mutation is associated with ocular anterior chamber dysgenesis. In a study of 55 cases with craniosynostosis syndromes, Tay et al [23] found a 35.5% prevalence of bilateral visual impairment and a 9.1% prevalence of unilateral visual impairment. The reported causes of visual impairment included amblyopia (16.7%), ametropia (25%), optic atrophy (16.7%), exposure keratopathy (4.2%), and infantile nystagmus syndrome (4.2%) [23]. Proptosis and strabismus are the hallmark ocular signs of Pfeiffer syndrome. Multidisciplinary management of ocular signs of Pfeiffer syndrome includes artificial tears, a lubricating ointment, eye care to prevent amblyopia, craniotomy for repair of craniosynostosis, and cosmetic reconstruction for facial dysmorphisms [24,25].

## Acknowledgments

This work was supported by research grants NSC-99-2628-B-195-001-MY3 and NSC-101-2314-B-195-011-MY3 from the National Science Council, Taiwan, and MMH-E-102-04 from Mackay Memorial Hospital, Taipei, Taiwan.

## References

- [1] Vogels A, Fryns JP. Pfeiffer syndrome. Orphanet J Rare Dis 2006;1:19.
- [2] Cohen Jr MM. Pfeiffer syndrome update, clinical subtypes, and guidelines for differential diagnosis. Am J Med Genet 1993;45:300–7.
- [3] Robin NH, Falk MJ, Haldeman-Englert CR. FGFR-related craniosynostosis syndromes. In: Pagon RA, Bird TD, Dolan CR, Stephens K, Adam MP, editors. GeneReviews™ [Internet]. Seattle, WA: University of Washington; 1993 [Updated 2011 Jun 7].
- [4] Oliveira NAJ, Alonso LG, Fangiello RD, Passos-Bueno MR. Further evidence of association between mutations in *FGFR2* and syndromic craniosynostosis with sacroccygeal eversion. Birth Defects Res A Clin Mol Teratol 2006;76:629–33.
- [5] Chen C-P, Lin S-P, Su Y-N, Chien S-C, Tsai F-J, Wang W. Craniosynostosis and congenital tracheal anomalies in an infant with Pfeiffer syndrome carrying the W290C *FGFR2* mutation. Genet Couns 2008;19:165–72.
- [6] Lai AHM, Tan Y-M, Law H-Y, Yeow VKL. A mutation in *FGFR2* in a child with Pfeiffer syndrome and a sacral appendage. Clin Dysmorphol 2008;17:73–4.
- [7] Stoler JM, Rosen H, Desai U, Mulliken JB, Meara JG, Rogers GF. Cleft palate in Pfeiffer syndrome. J Craniofac Surg 2009;20:1375–7.
- [8] Chen C-P, Su Y-N, Hsu C-Y, Lin P-Y, Tsai F-J, Chern S-R, et al. Second-trimester molecular prenatal diagnosis of sporadic Apert syndrome following sonographic findings of mild ventriculomegaly and clenched hands mimicking trisomy 18. Taiwan J Obstet Gynecol 2010;49:129–32.
- [9] Tartaglia M, Valeri S, Velardi F, Di Rocco C, Battaglia PA. Trp290Cys mutation in exon IIIa of the fibroblast growth factor receptor 2 (*FGFR2*) gene is associated with Pfeiffer syndrome. Hum Genet 1997;99:602–6.
- [10] Schaefer F, Anderson C, Can B, Say B. Novel mutation in the *FGFR2* gene at the same codon as the Crouzon syndrome mutations in a severe Pfeiffer syndrome type 2 case. Am J Med Genet 1998;75:252–5.
- [11] Ariga H, Endo Y, Ujiie N, Ishii T, Ishibashi N, Fujita T, et al. Trp290Cys mutation of the *FGFR2* gene in a patient with severe Pfeiffer syndrome type 2. Pediatr Int 2001;43:293–5.
- [12] Shotelersuk V, Ittiwut C, Srivuthana S, Mahatumarat C, Lerdlum S, Wacharasindhu S. Distinct craniofacial-skeletal-dermatological dysplasia in a patient with W290C mutation in *FGFR2*. Am J Med Genet 2002;113:4–8.
- [13] Zackai EH, McDonald-McGinn DM, Stolle C, Huff DS. Craniosynostosis with tracheal sleeve: a patient with Pfeiffer syndrome, tracheal sleeve and additional malformations in whom an *FGFR2* mutation was found. Clin Dysmorphol 2003;12:209.
- [14] Hockstein NG, McDonald-McGinn D, Zackai E, Bartlett S, Huff DS, Jacobs IN. Tracheal anomalies in Pfeiffer syndrome. Arch Otolaryngol Head Neck Surg 2004;130:1298–302.
- [15] Nazzaro A, Della Monica M, Lonardo F, Di Blasi A, Baffico M, Baldi M, et al. Prenatal ultrasound diagnosis of a case of Pfeiffer syndrome without cloverleaf skull and review of the literature. Prenat Diagn 2004;24:918–22.
- [16] Lajeunie E, Heuertz S, El Ghouzzi V, Martinovic J, Renier D, Le Merrer M, et al. Mutation screening in patients with syndromic craniosynostoses indicates that a limited number of recurrent *FGFR2* mutations accounts for severe forms of Pfeiffer syndrome. Eur J Hum Genet 2006;14:289–98.
- [17] Koga H, Suga N, Nakamoto T, Tanaka K, Takahashi N. Clinical expression in Pfeiffer syndrome type 2 and 3: surveillance in Japan. Am J Med Genet 2012;158A:2506–10.
- [18] Gonzales M, Heuertz S, Martinovic J, Delahaye S, Bazin A, Loget P, et al. Vertebral anomalies and cartilaginous tracheal sleeve in three patients with Pfeiffer syndrome carrying the S351C *FGFR2* mutation. Clin Genet 2005;68:179–81.
- [19] Akai T, Yamamoto K, Iizuka H, Kawakami S, Yoshida J, Kakinuma H, et al. Syndromic craniosynostosis with elbow joint contracture. Pediatr Neurosurg 2006;42:108–12.

- [20] Stevens CA, Roeder ER. Ser351Cys mutation in the fibroblast growth factor receptor 2 gene results in severe Pfeiffer syndrome. *Clin Dysmorphol* 2006;15:187–8.
- [21] Ranger A, Al-Hayek A, Matic D. Chiari type 1 malformation in an infant with type 2 Pfeiffer syndrome: further evidence of acquired pathogenesis. *J Craniofac Surg* 2010;21:427–31.
- [22] Okajima K, Robinson LK, Hart MA, Abuelo DN, Cowan LS, Hasegawa T, et al. Ocular anterior chamber dysgenesis in craniosynostosis syndromes with a fibroblast growth factor receptor 2 mutation. *Am J Med Genet* 1999;85:160–70.
- [23] Tay T, Martin F, Rowe N, Johnson K, Poole M, Tan K, et al. Prevalence and causes of visual impairment in craniosynostotic syndromes. *Clin Experiment Ophthalmol* 2006;34:434–40.
- [24] Harb E, Kran B. Pfeiffer syndrome: systemic and ocular implications. *Optometry* 2005;76:352–62.
- [25] Iannetti G, Ramieri V, Pagnoni M, Fadda MT, Cascone P. LeFort III external midface distraction: surgical outcomes and skeletal stability. *J Craniofac Surg* 2012;23:896–900.